

*American Handbook of Psychiatry*

# PHYSIOLOGICAL SOCIOLOGY:

*Endocrine Correlates of Status Behaviors*

Patricia R. Barchas

Jack D. Barchas

# **Physiological Sociology:**

## **Endocrine Correlates of Status Behaviors**

**Patricia R. Barchas and Jack D. Barchas**

e-Book 2015 International Psychotherapy Institute

From *American Handbook of Psychiatry: Volume 6* edited by Silvano Arieti

Copyright © 1975 by Basic Books

All Rights Reserved

Created in the United States of America

## Table of Contents

### PHYSIOLOGICAL SOCIOLOGY: ENDOCRINE CORRELATES OF STATUS

#### BEHAVIORS

#### Introduction

#### Status as a Social Behavior

#### Some Endocrine Correlates of Status Related Behavior

#### Psychiatric Processes and Physiological Sociology

#### Concluding Remarks

#### Bibliography

# PHYSIOLOGICAL SOCIOLOGY: ENDOCRINE CORRELATES OF STATUS BEHAVIORS<sup>1</sup>

## Introduction

Intermittent concern, variously expressed, with the relationship of individual physiology to social situations has been with social psychology at least since that field borrowed the concept of homeostasis from biology. The newer balance theories in the social sciences share with earlier views the central notion that disruption of the system produces effects on the social, psychological, and physiological levels and that the system will tend to attempt to reduce that tension by reestablishing balance (homeostasis). We would like to present some of the relationships and considerations in this chapter that have been made between a basic social process and endocrine measures that have been related to homeostasis and behavior.

Sociology is concerned in part with the ways in which group structure influences individual behavior. Status is viewed as one of the basic structural processes of social behavior in humans and other animals, although expressed differently in different species. The importance of status as a concept for social behavior would parallel the importance of the concept of the utilization of glucose for biochemistry. Although not generally recognized by persons with training in "psychological" areas, concepts such as status are the heart of investigation for social scientists involved in the study of

interactional processes. We would like to examine some aspects of status as an example of a basic social behavior and present some of the correlations that have been made between status behavior and endocrine behavior.

We are presenting this system of organization for heuristic reasons: it permits a juxtaposition of two stances of inquiry, each of which seems relevant to psychiatry and can be made relevant to each other. However, utilizing such a system for some purposes does not preclude recognizing the value of other ways of inquiring into the relationships of physiology to behavior.

The physiological responses with which we will be concerned in relation to status behavior will emphasize hormonal patterns involving the adrenal-cortical steroids, adrenal-medullary and gonadal hormones, and brain neuro-regulators. Each of these endocrine or neuro-regulatory systems has, at some time, been considered as one of the body's systems of homeostatic mechanisms; each of the systems is activated or depressed under differing stimulus conditions; and each remains in effect for differing lengths of time, and influences other systems of response differentially. Individual variation is great, and, the data to date can in no way be applied as an explanation to any given single event. The knowledge that has been accumulated represents promise, but the needs for further information in the systems directly mentioned, and in systems which have not been studied, is

very great.

We believe that investigation of these areas has great relevance for psychiatry. A variety of illnesses has been identified in which “psychological” factors may play a major role. These range from forms of allergy, hypertension and dermatoses to disorders of the gastrointestinal tract among others. For any processes that are so construed, social processes such as status, dominance, and role relationships and their complexities may be crucial parts of what has been considered “psychological.” In the past, psychological processes in such illnesses have frequently been considered in terms of individual psychology rather than in terms of structural processes involving the individuals. In this paper we will deal exclusively with such structural explanations, although we are also convinced of the great importance of individual psychological processes.

We will present a general description of the concept of role and status. We will then proceed to a more detailed description of status behavior in a variety of human and nonhuman primate studies, studies that suggest powerful evolutionary aspects in this behavior. Some of the literature relating status behavior to endocrine function will be presented. Finally, we will consider aspects of the study of physiological sociology in terms of the relation of the physiological and behavioral events and the manner in which they may interact. We will take up some of the possible relations of these

materials to psychiatric illness.

## Status as a Social Behavior

### Some General Aspects of Role Therapy and Status in Human Groups

“Group dynamics” insofar as they are repeated group to group, independent of particular persons, may be considered as structural in nature. Groups themselves may be voluntary or involuntary, formal or informal, open or closed. But, without a structure (patterned relationships) no aggregate of people may be considered a group. One way of looking at group behavior, which permits us to relate it to physiology, is in terms of roles. Some of the basic ideas used by role theorists are: persons are the only actors in the social system and these persons occupy positions (statuses). To these positions are attached, to a greater or lesser degree, expectations about behavior. These expectations, of rights and obligations associated with the positions people take, constitute a role. There is room for individual style in carrying out rights and obligations. This is not part of the role. A position is delineated by shared expectations, and a role is delineated by behavior that reflects positional expectations. (It should be noted that the position—status—as well as the way in which a role is enacted, may be evaluated by self or others either positively or negatively. We will not be dealing with such evaluations, even though viewed as reward and punishment, we believe they bear upon status



and physiological response to status.)

Relevant to role behavior are socialization into roles and maturational limitations on the abilities of actors to carry out roles. This area of development is of extraordinary importance and is essentially unexplored.

Any person carries about with him any number of role possibilities. Dependent upon the social situation, one or another may be activated. Each individual must be able to move from role to role, sometimes to juggle several simultaneously, and sometimes to find that role requirements are either ambiguous or in conflict with self-concept. Any and all of these can be termed role conflict; as such, for both literature and social psychology, as well as for psychiatry, they produce the stuff out of which the human drama is written.

Tension, its accompanying pains and consequent psychological or socially potential disruptions; inappropriate reduction mechanisms, again psychological or social in nature; and distortion of either self-concept or the social structure may result from role conflicts. It is not surprising that we may find physiological correlates of such powerful and ubiquitous human phenomena. Because of rapid social change, the roles for which children prepare, and the coping mechanisms they learn, may not meet the demands of their changing adult life situation.

Intrinsic to the idea of roles is the notion of reciprocity. For each status

or position, there is a counter-status; e.g., teacher-pupil; mother-child; friend-friend; employee-employer. Clearly, these may be hierarchically or horizontally arranged. Nevertheless, the rights and obligations that define the relationship are bidirectional, although not necessarily of equal intensity or frequency of emission. There are cases (probably most available to the therapist and difficult to document as to validity) in which the reciprocal roles for the individual in question are purely symbolic, as is the actor to whom the role is attached. Thus, we may have images of absent figures influencing our actions. These figures are sometimes constructed from childhood experiences, fantasy, literature, or mythology. For sociological studies, one would be concerned primarily with simultaneously present actors who simultaneously activate reciprocal roles.

One way to view notions of role is as a principle of organization that is meaningful, consciously or unconsciously, both to the individual actor and to the observer. We may view culturally held norms transmitted to the individual as general and specific attitudes. These attitudes are further organized around societal statuses or positions, and are emitted as kinds of behavior.

Remembering that the individual is more than the sum of his statuses and roles, we nevertheless may claim that much social behavior, if not all, is partially governed by role considerations. The specifications for behavior are

more or less rigid and vary in degrees of clarity. Usually, individuals carry out their roles with a minimal amount of conflict and uncertainty. However, whenever a social situation activates two roles simultaneously, there is the possibility for conflict. In situations in which role requirements are highly specified, there is an increased possibility that required behavior will be at odds with the individual's conception of himself, or even with higher order role commitments as perceived by the individual. Plainly, situations in which role requirements are not clear can be stressful. Plainly, too, there is great individual variation in both the mode of dealing with, and the degree of, felt stress involved in these situations.

Areas of inquiry that may be organized in terms of role considerations—although they may equally well stand-alone—include power relations, coalition formation, leadership phenomenon, and affiliative bonding.

## **Ubiquity of Status**

We are working from the point of view that man is a biological organism, similar to other organisms, especially other primates, and that his biology is a determinant of the range of his behavior, social as well as physical. His particularly human characteristics we take to be derived from the cortex: these include plasticity, use of experience, use of symbols and language, and time sense.

Rejecting “social Darwinism,” we nonetheless believe that social structures and social processes that are found to be most widely distributed in small human groups will be found in other primates as well. Should this be true, one could project to underlying mechanisms, perhaps biological, that partially determine such processes. Casting our ideas in an evolutionary framework, we believe there is a substrate of social forms and processes, necessarily limited to small group interaction, that is common to the primates, including man.<sup>o</sup> We tentatively agree with Colter Rule who suggests that some kinds of social behavior, such as dominance “are as fundamental to human existence as such short rhythm phenomena as respiration and the heartbeat: intermediate (circadian) rhythms like sleeping, waking, eating, drinking; and longer rhythms like sexuality and seasonal changes.” We think it even more likely that some social processes involving structure are basic—such as some status processes, affiliative bonding of several types, and some depressive reactions.

The ubiquity of status orderings in small groups has been repeatedly confirmed by observers of mammalian behavior. Whatever the nature of the population, status is generally thought to influence such social behavior as leadership, coalitions, aggression, priority of access to desired objects, intragroup conflict, conformity, and group structure.” Given these relationships and the current proclivity to compare human with nonhuman behavior—and to seek animal models for studying human social phenomena

—it seems reasonable to explore the status phenomenon in a variety of mammals, so that methods and results can be compared with results from human studies. This is particularly true if there continues to be an increased tendency among behavioral scientists to assume that basic social behavior is in part biologically determined, and that the similarity in the biology of nonhuman organisms and humans may dictate similar social processes at the most elementary level, although the specific social patterns and their meanings may rest on other foundations.

This view does not revive or take sides in the nature-nurture argument. It does, however, have evolution as its framework. It argues that the brain, like other organs, retains substrata, and that these substrata partially govern the most basic kinds of behavior, such as learning, emotional behaviors such as depressive reactions, and some social interactions. While the effect of both further evolution and the impact of language and culture is not denied, this view does present the possibility that some of man's social responses are more adapted to past patterns of small group interaction, given the lag between culture and man's biologic adaptation. It also suggests that the most basic social responses may be studied in nonhuman populations with the same advantages and disadvantages as are found, for example, in the study of learning among nonhuman primates.

### **Approaches to Status and Findings in Humans**

Status is, in general, an ordering of individuals by which rights, privileges, and responsibilities are distributed. The power and prestige order of a group is often considered synonymous with the status ordering. Status is a ubiquitous phenomenon in human groups. Even gross knowledge of status positions explains and predicts many social behaviors. A description of a group typically uses status to order observations.

There are three aspects to the phenomenon of status emergence: status orderings form, are maintained, and change. We will review the literature dealing with status formation, omitting the portion dealing with maintenance and change. Most studies of status formation focus on ad hoc, freely interacting task groups. (By task group, we mean any small group brought together for the purpose of achieving a mutual goal through mutual efforts.)

In this context, it has been found that diffuse status characteristics, age, sex, race, which individuals bring to the group, are likely to determine the status ordering within the group. Recently, Fisek and Berger presented material extending this approach to include any factor that discriminates among the actors, such as occupation, but not including personality and socialization variables.

We note that in such groups, even when diffuse characteristics are held constant, a status ordering emerges. The Balesian tradition has yielded most

of the data on such groups. The status order in freely interacting task groups forms quickly, in as little as forty-five minutes and it is stable. Working within this Balesian tradition, Fisek found that the differentiation occurred rapidly.

Two routes to status formation have been studied, one in which the status order develops out of interaction, and one in which the status order precedes apparent interaction. The work of Berger et al. may be taken as an exploration of the first route of formation, although it clearly deals with other matters as well. In this approach, both formation and maintenance of the status or power and prestige order are systematically linked to the task through performance evaluations and consequent expectation states.

The Berger model does not account for those groups in which the differentiation has taken place prior to interaction, particularly those cases in which the observed initial order is stable over time. The simplest approach to explaining initially differentiated groups may be to look for variables that might operate in ways similar to diffuse status characteristics. Personality variables or socialization variables are often examples, both expressing themselves in “style,” the cues for which could be extremely muted—carriage, posture, propensity for eye contact. If one were to predict which individual was to occupy a given position, one would probably need to take into account such variables systematically. The results may be culture bound, and, in addition, depend not only upon the background of the actor but upon the

perceptions of other actors.

Status as a process is especially suited to comparative study. As a phenomenon, status has received attention from sociologists of human behavior for many years, and there is agreement that status is an important determinant of human behavior. A large body of descriptive, experimental, and theoretical literature has arisen concerning status. Those observations which have been made on established ongoing groups suggest that status differentiation occurs universally, and that status differentiations correlate highly with performance in a group.

Thus, status differentiation apparently occurs universally in human groups, and status can be observed in both ongoing and experimental groups. Further, when members of a group have a collective task, they evolve patterns of interaction that clearly reflect differences in power and prestige among the members of the group, even when they are strangers at the onset of observation and are matched for external status characteristics.

### **Findings on Status in Non-Human Primates**

Similarly, observers of nonhuman mammals which live in groups have found some form of status differentiation to be generally present. We restrict our comments in this section to nonhuman primates. Starting with Carpenter, field observers have used the notion of status to order their observations. The



term status has frequently meant dominance and has connoted priority of access.

Field workers and observers in semi-natural habitats have described aspects of group structure for many species, cataloguing and describing behavior in context. These studies generally have been done either from a straight analogic approach or from an evolutionary approach to behavior. It has been assumed that there may be common forms of social behavior across related species, if not common meaning for each form. From these studies comes the suggestion that there are intriguing similarities in the organization of social behavior across primate species, the similarities becoming greater as higher levels of abstraction are used in behavioral categories.

Observational categories have been developed that are apparently reliable across observers. There has developed a concurrence regarding the “meaning” of some kinds of behavior, relative to slightly higher order concepts, such as dominance and deference. There is seldom a report in the primate literature in which some reference is not made to a dominance order, if not a status order, among the group.

In general, the high-status animal has freedom of movement, other animals making way for the high-status animal who may take the desired objects, such as food, from the other animals.’ Data have been interpreted to

mean that disruption of the status ordering causes instability in the group and that such variables as extent of group territory correlate with the temperament of the highest status individual in the group. It has been suggested that the sleep of the high-status animal may be deeper and less disturbed than that of low-status animals. It has been found that the later status position of an infant is influenced by the present status of the mother. Further, it is suggested that types of mothering are affected by the status of the mother. In general, it has been found that adult males have higher status than adult females who, in turn, have higher status than young animals, although there may be considerable overlap in these orderings.

Most observations of nonhuman primate groups have been made of single groups of mixed age and sex. There is consensus among observers that status relationships partially order behavior. It has generally been assumed that these relationships are the product of long term socialization in stable groups, and that the advent of status relations developing in new groups can be attributed to external differentiating characteristics in the animals, such as sex, size, and weight. Goodall and Hamburg suggest, on the basis of observations of free-ranging chimpanzees, that motivation and “technical ingenuity” may be factors in an individual’s rise in dominance.

Status differentiation has been found to be stable in naturally occurring groups, although there are clearly conditions under which status shifts occur,

these being related to growth and development as well as to cyclical and other biological changes. Bernstein and Mason, reporting on a newly established, mixed group, state that status relationships formed within the first hour of interaction were stable for seventy-five subsequent days.

Focusing upon the dominance aspect of status in groups of mixed status, Bernstein and Mason have demonstrated that dominance relations in mixed groups were quickly formed. Miller and Murphy report that such relations are stable, and Warren and Maroney and Kawai have demonstrated them to be highly resistant to change. These features have been attributed to (1) external status differences, or (2) long-term socialization of the individuals with each other. However, in a study by Barchas these two attributions were ruled out as both were controlled for. Thus, animals with fully developed social repertoires have the capacity to form status orderings out of interactions which are independent of broad differentiating cues and of interaction with known individuals.

In mixed groups, the fully adult male dominates females and younger animals, the larger animal the small, the animal in good condition the animal in poor condition. Most observations are upon groups of varying statuses.

Clearly, both human and nonhuman primates in field and laboratory settings have been observed to exhibit status orderings. As has been stated, in

humans the basic finding about status orders comes from Bales and his associates, “. . . marked inequalities develop over time in the rate at which members are observed to initiate interaction [and] those who initiate action most frequently tend to be ranked highest on the criteria of ‘best ideas’ guidance and tend to receive actions from others at the highest rate.” This occurrence has been further documented in a variety of contexts and its stability established with little question. For specific information on contexts other than the original Bales study, see Berger et al., Investigation into the phenomenon as a process and on a theoretical and experimental level has been most vigorously pursued by a group of Stanford sociologists.

## **Some Endocrine Correlates of Status Related Behavior**

### **Adrenal Cortical Functions**

Until recently, the relationship between status and endocrine function has been investigated primarily in rodents. Such studies have tended to focus on the relationship of adrenal cortical function to dominance (one dimension of status) and have yielded fairly consistent results. Thus, the studies of Barnett regarding fighting behavior in wild rats suggested, in a dramatic fashion, that the adrenal cortical hormones may be markedly depleted following fighting in the subordinate animal, but not in the dominant animal, although both appeared to have had an equal overt stress. In a study of male

mice derived from a wild strain, it was noted that there was an inverse correlation of the adrenal weight and social rank in animals having had social interaction over a ten-day period; high ranking animals have less adrenal weight than low ranking mice. Such findings may be related to physiological changes occurring in crowded conditions, where it has been found that in rodents increased group density results in increased adrenal weight, but decreased body and testicular weights. Isolated male mice display lower resting levels of plasma corticosterone and a decreased response to stress. If mice are maintained in a paired situation for several weeks, the subordinate mice are characterized by higher adrenal weights and lower gonadal weights; however, if tested in a single trial against a new non-aggressive antagonist, the subordinate animals behave within the same range of behavior as the dominant animals. The author notes that it is not surprising that the later behavior is independent of the earlier behavior, since the behavior is part of an established hierarchical response and the response to an unfamiliar individual should be independent.

Direct measures of adrenal activity by measurement of plasma corticosterone as an indicator of adrenal-cortical activity have been obtained. They confirmed the earlier impressions based upon adrenal weight: dominant mice had significantly lower levels of corticosterone than subordinate animals. The causal nature of the relationship has not been established.

Primate studies have provided a quite different view. Although such studies are difficult, the work of Leshner and Candland is particularly interesting in the suggestion that higher status is associated with increased adrenal-cortical activity. Their work utilized assay of urinary 17-hydroxycorticosteroids with correlation to dominance ranking in an ongoing group of squirrel monkeys. Such a view is confirmed through adrenal weight measures, though without kinetic endocrine measures, in work with crab-eating monkeys.

An interesting but difficult investigation to interpret in light of the preceding studies dealing with primates, comes from the work of Sassenrath who studied the response in rhesus to adrenocorticotrophic hormone (ACTH) on urinary steroids collected outside of the behavioral situation. In that study, the dominant animal has the lowest secretion of 17-hydroxycorticosteroids in response to ACTH. The interpretation would be that the response to ACTH is a test of the adrenal size, and that the lower the response, the less the activity of the adrenal. Although the interpretation is complicated, the results suggest powerful influences of social behavior on pituitary-adrenal function.

If one accepts the Leshner and Candland and Hayama data, there are profound differences between the rodent and primate in regard to pituitary-adrenal function. The differences are viewed by Leshner and Candland as due to the different styles of ongoing interaction. Thus, in the rodent, dominance

is established by direct physical attack, while in the primates that have been studied the dominance is settled by ritualized displays. These processes may relate to varying forms of aggressive interaction which have differing underlying endocrine and neurochemical mechanisms.

Studies dealing with endocrine measures in primates are difficult because of the problems associated with obtaining adequate or appropriate samples of blood in relation to ongoing behavior. Thus, frequently the collection of the compound of interest is accomplished under conditions that are different from those involved in the status situation or social behavior situation per se. One could anticipate that for some kinds of behavior, there may be marked differences between the social situation and the collection situation in terms of endocrine response.

### **Gonadal Function in Males**

Only very recently have procedures been developed that enable direct investigation of testosterone in relation to social behavior. Not only have the chemical assays required development, but the behavioral procedures for the collection of blood samples have also required development, so as to be able to obtain the samples very quickly in adapted animals. As a consequence of such development, important studies in which kinds of behavior could be determined with good reliability in relation to the endocrine measure have

become possible. From studies involving thirty-four animals in a one-third acre compound, it was established that dominance rank was positively correlated with testosterone concentration. The animals in the highest quartile had significantly lower levels of testosterone than those in the lower quartile, although there were no significant differences between animals in the second, third, or fourth quartiles. Aggression also correlated with testosterone, but the correlation of high submissiveness and low testosterone was not high. The complexity of the data is shown in the fact that the relationships are not always simply related. Thus, the most aggressive animals are not always the most dominant, and the most dominant animal need not have the highest testosterone. The studies led the investigators to a series of investigations of the relationship of behavior to endocrine changes, and an attempt to determine whether testosterone changes were preceded by changes in dominance. The data to investigate this possibility was obtained by using smaller groups and allowing animals to take a dominant position, in which case plasma testosterone increased; if the dominance position sharply decreased, the plasma testosterone decreased. These results are also of interest in light of the findings that plasma testosterone in humans is decreased by stress.

## **Adrenal-Medullary Function**

Relations of adrenal-medullary secretion to behavioral processes



related to social behavior has proceeded upon two fronts. The first has been the area of measurement of adrenal-medullary hormones, epinephrine (adrenaline), and norepinephrine (noradrenaline), generally utilizing urinary samples. The second, in animals, has involved studies of the enzymes that form the catecholamines in the adrenal, as well as measurement of the catecholamines, with determination of changes associated with behavioral states. The rapid changes that can occur in the adrenal-medullary system and the powerful psychological effects of epinephrine makes this system a particularly interesting one for investigation.

Several studies suggest that social behavior may alter the enzymatic processes involved in the formation of the catecholamines. Among those areas of investigation are studies dealing with social isolation and group housing, as exemplified by the studies of Welch and Welch. They found decreased levels of catecholamines in the brain and adrenal of group-housed mice when contrasted with individually housed animals. Although the results of such studies may depend upon the species used, it is clear that differential housing can markedly alter catecholamine mechanisms. More directly related to social interaction have been the investigations represented by the collaboration of Henry and Axelrod and their colleagues. They have utilized a variety of mutual interaction situations including interconnected cages, a technique which leads to confrontations and severe social stimulation. In animals exposed to such severe social interaction, there were increases in the

enzymes that form catecholamines in the adrenal, as well as increases in the levels of the hormones themselves. A decrease in the enzymes was noted in isolated animals. Preliminary studies have suggested that dominant animals may have lower levels of the enzymes that form catecholamines than the subordinate animals have. Many of the studies of the investigators utilized long-time bases (in some of the studies, periods as long as six months were used) and it will be of interest to see if changes associated with these and other social interaction paradigms can be demonstrated with short-time span interactions. Rapid changes in the enzymes that form catecholamines have been noted to occur within a few hours of certain forms of stress, which suggests the possibility of investigations involving behavioral states with short-time parameters in the appropriate social situation.

Studies of adrenal-medullary function in humans have concentrated on determination of epinephrine and norepinephrine in urine. This has been necessary because of the difficulty in measuring the catecholamines in blood samples given the limitations of current assay procedures. Thus, samples that are assayed represent a pooling of many time-dependent processes, usually over a period of several hours, which is unfortunate, yet makes the fact that there are positive results even more tempting for future analysis with multiple time points. Ultimately, one would expect that assessment of adrenal-medullary function in humans would involve determinations at close time intervals by means of new techniques, which are now being developed,

such as mass fragmentography.

Several investigators have demonstrated powerful effects of psychosocial interaction variables on adrenal-medullary secretion. Many of the studies have demonstrated that a variety of behavioral situations that involve tense, anxious but passive emotional displays are associated with elevated epinephrine output. This can be said of certain novel or distressing situations as well. A variety of psychological states have been investigated in a pioneering series of studies conducted over many years by Levi and brought together in a recent monograph. Urinary catecholamines were found to increase in a wide variety of arousing situations, such as viewing films with both "pleasant" and "unpleasant" aspects. Bland materials reduced the levels of excretion. Psychosocial stimuli were found to alter catecholamine secretion, depending upon the stimuli and on the starting point of the organism, without any simple relationships between anxiety and epinephrine secretion or aggression and norepinephrine secretion.

Frankenhaeuser has conducted a series of careful investigations relating cognitive and emotional patterns with endocrine secretion. Among the most interesting studies has been a series of investigations in children. These, while not specifically concerned with status, suggest powerful social interrelationships that may bear on status and power relationships. For example, there was a positive correlation between the norepinephrine

secretion of mother and son, but not of father and son. There appeared to be a significant positive correlation between the mother's adrenaline output and the frequency with which the fathers punished their children. Such studies of interactional and developmental processes, particularly when combined with social processes, suggest outlines for the development of a new research area.

A means of studying the relationship of social processes to endocrine processes has emerged from investigations of free fatty acids and social behavioral states. Free fatty-acid levels are believed to correlate highly with sympathetic activation of which adrenal-medullary activity would be an important component. In a series of studies characterized by the use of rigorous behavioral parameters, as well as rigorous chemical measures, Back and Bogdonoff have demonstrated correlations between chemical measures and behavioral measures of conformity and leadership in test situations in which these parameters could be directly controlled. Using changes in free fatty acids as an indicator, the authors of these pioneering studies note:

... if the social situation is perceived simply as a background to individual achievement, the dominant variable will be the potential for individual achievement and its meaning. In this situation, pressure to conform and pressure to assume leadership may be viewed as arousing stimuli, and the individual may seek to avoid these situations. If, however, the group relationship has the dominant meaning, the performance of the task may be seen in terms of the group interaction, and then deviation from the group norm becomes the arousing condition. Conforming behavior is then attended by decreased arousal, [p. 41 ]

A direct study of the relationship between social status processes and catecholamine secretion has just been completed by Barchas and Barchas. In this study, individuals of equal external status (education, age, race, sex) were brought together, and they interacted for a one-hour period. The status differential that was established between the individuals through interaction was assessed by observers and by questionnaires for the subjects. Urinary catecholamines (epinephrine, norepinephrine, and dopamine) were obtained before and after the session. There was no relationship between the urinary catechols taken before the interaction and the acquired status, but there -was a pattern of association between acquired status and the urinary catechols sampled after interaction.

### **Brain Serotonin**

Biogenic amines in the brain have been studied in terms of their relation to several kinds of behavior, including severe psychiatric disorders. Serotonin, an indoleamine, has been postulated to be a possible transmitter and to act as a regulator of neuronal function. The compound has been linked to specific forms of behavior, but only limited studies related to social behavior have been conducted. A major procedure for investigating the effects of changes in brain serotonin involves use of parachlorophenylalanine (PCPA) which inhibits the formation of serotonin, although it may also act on other chemical systems. Studies in which PCPA has been administered to

primates have been very few and suggest decreased activity. Boelkins administered the drug to an ongoing group of three crab-eating macaques (*Macaca fascicularis*) and observed that the number of kinds of social behavior decreased, but that the time spent in social huddling increased, with no change in aggressive or other kinds of behavior.

Maas et al. in studies with two groups of *Macaca speciosa* observed no changes in aggressive or submissive gestures, attacks, or hetero- and auto-grooming.

In each of these studies, the degree to which brain serotonin mechanisms have been interfered with is open to question, and larger doses of the inhibitor of serotonin formation caused the animals to appear ill, possibly due to peripheral effects. Thus, the role of serotonin in primate social behavior deserves further investigation.

## **Brain Catecholamines**

Particular attention has focused on catecholamines as they relate to a variety of behavior. By the use of specific inhibitors of the formation of catecholamines, studies have been performed suggesting important roles for catecholamines in social behavior. In a series of papers, the group associated with Maas' has investigated such processes in studies of macaque monkeys. In one set of investigations, the drug  $\alpha$ -methyltyrosine, an inhibitor in the first

step of catecholamine synthesis, was administered to some of the animals in a group-living situation, with monitoring of social behavior. It was found that the animals had a marked decrease in social initiation, including grooming, threats, and attack, although they responded to other animals. There also may be dominance shifts: an animal who was dominant did not have a change in social dominance, while two other animals had a decrease in dominance. In another study, a different drug, 6-hydroxydopamine, which destroys many of the neurons that contain norepinephrine in the central nervous system, was administered to free-ranging macaques. The treated animals were more peripheral to their social group, exhibited decreased social behavior, initiated fewer threats, less social grooming, and fewer social initiatives than the control animals.

The results of the Maas group demonstrate powerful effects of central catecholamines in social processes in nonhuman primates. Such studies would suggest that certain forms of social behavior may be mediated by catecholamine-containing neurons and be altered by drugs that alter catecholamine mechanisms. The processes that have been studied are related to basic social processes previously shown to be related in similar ways in nonhuman and human primates. The findings that central catecholamine systems are profoundly involved in nonhuman primate social behavior raises questions as to the role of catechols in humans.

## Psychiatric Processes and Physiological Sociology

### Some Aspects of Sociopathic Behavior

A number of psychiatric processes are observable in which there is clinical reason to believe that profound disturbances of sociological processes, including status, dominance relations, and affiliative bonding exist. One psychiatric illness that manifests itself in social structural relationships in a profound way is sociopathic behavior.

By the very nature of the sociopathic individual, persons with the disorder display altered social behavior. The question as to whether in terms of their physiological responses such individuals respond differentially when compared to normal people has been raised by several investigators. Some evidence suggests that there are such differences.

Learning variables are among the factors that have been investigated. The studies of Hare and others have suggested that classical conditioning of autonomic responses and avoidance learning may be impaired. For example, it has been found that psychopathic criminals acquire conditioned electrodermal responses more slowly, that the formed responses are of lesser magnitude and are more quickly extinguished, and that, once acquired, the responses are generalized to a lesser degree than in non-psychopathic criminals. These studies and others suggest that psychopaths may have



normal attention to stimuli, but a decreased anticipatory fear.

Particularly relevant from the standpoint of this review are a series of provocative studies dealing with the response to an injection of adrenaline by psychopaths when compared to other populations. The literature has been quite contradictory, but does strongly suggest that there are differences.' The experiments were performed under different conditions, including degree of ongoing activation. In the studies of Hare, which were able to take advantage of the earlier findings and to work with a carefully selected population, it was found that the change in skin conductance following the injection of adrenaline was smaller in a psychopathic group compared to a criminal non-psychopathic group. In that study, differences in heart rate that had been noted in the earlier studies, with quite different subject-selection procedures and activities, were not found. Questions as to the hyper- or hyposensitivity of aspects of the autonomic nervous system have been raised by the various studies, although conclusions would require considerably more investigation.

Another side of the coin remains to be investigated—the effects of stresses on direct endocrine release. Such studies would involve, for example, in relation to the previous literature, determination of release of epinephrine and norepinephrine and the enzyme that converts dopamine to norepinephrine (dopamine- $\beta$ -hydroxylase), which can be measured in the peripheral blood. Ideally, diurnal rhythms and the response to stress, and

particularly the response to emotional conditioning, would be obtained. Studies of central brain mechanisms are clearly indicated.

The literature dealing with psychopathic behavior raises the question as to whether this form of social behavior, which involves a psychiatric diagnosis, may not have powerful biological aspects. One could imagine either biochemical differences based on genetic factors or on early experience. In either event, it is quite likely that psychopathic illness may reflect interactions between social behavior and biochemical processes<sup>2</sup>

Frequently role enactment requires simultaneous action or interwoven action between two or more individuals. Discrepancies in perceptions of roles may lead to group stresses. Most frequently in this situation there is a mutual adjustment of expectations. When there is not, we may have such phenomenon as schism and skew in the family and scapegoating (here seen as a mode of releasing group tensions regarding roles to the detriment of the individuals involved). Similarly, the phenomenon of the double bind seen from the points of view of the sender and receiver of messages may be partially visualized as inadequate internalizations and expressions of role relations. Such social behavior could be expected to have profound effects on the types of parameters considered in this chapter. An additional example of the application of role theory and status processes to psychiatric illness could be made in the case of depressive illness.

## **Toward a Physiological Sociology**

It is difficult at this early point in the development of physiological sociology to give an adequate general formulation of the potential relations between physiological events and sociological behavior. At this stage, it seems reasonable to assume that biochemical and sociological processes are intimately related. For example, biochemical processes may affect activity levels, emotional tones, and the susceptibility to stress. Sociological processes may set in motion processes that influence biochemical mechanisms; for example, changing the levels of particular compounds by producing shifts between pathways, altering utilization of compounds, and inducing enzymatic changes. Biochemical events may profoundly alter the ability of the organism to respond to its environment.

If one visualizes a relationship between biochemical and social-structural events and assumes changes in response to a sociological event, then such changes become very important in relation to the length of time that they persist and the manner in which environmental and genetic factors interrelate. To account for long-term emotional behavior and alter-ability of behavior, one would have to assume alterability in underlying chemical events. In the simplest model, we would assume that sociological events affect the body chemistry and that the chemical change in turn affects the future sociological events. Thus, with particular genetic predispositions, chemical

changes that are long-term in nature could, in effect, “lock in” certain psychological sets.

Different genetic strains have already been demonstrated to have a considerable variation in terms of the steroid hormones produced by the adrenal cortex and thyroid hormones. In a number of illnesses it has been demonstrated that there is a genetic difference in formation of adrenal-cortical hormones; several of these illnesses lead to marked behavior change.

Strong evidence now suggests that some of the other biochemical systems we have been concerned with in this chapter are also under genetic control. Thus, the levels of the enzymes that synthesize catecholamines (epinephrine, norepinephrine, and dopamine) vary in different inbred strains. The degree to which the level of the enzymes can be altered by stress and even the type of stresses to which there are responses varies in different inbred strains.

An extensive listing could be made of how possible genetic differences in a system such as that involved in catecholamine production or utilization, both in the adrenal or in the brain, might affect social behavior. It has been shown that in response to stress differing amounts of adrenal-cortical steroids are released in different genetic strains. It is easy to imagine that in response to stress two individuals might send the same number of nerve

impulses to the adrenal medulla, but, because of a genetic difference, one individual might form differential amounts of catechols, release differential amounts, metabolize the catechols at different rates, or have differential passage across the blood-brain barrier. If any of these possibilities were to occur, clearly there could be behavioral changes when the adrenaline reached various target organs, including the brain.

Other possibilities to be investigated include the presence of minor or abnormal pathways in adrenal-catechol metabolism, including the physiological controls on those pathways, and how they might be altered by social stress. Analogous factors could be relevant not only to adrenal catecholamines as hormones related to behavior but also to brain catecholamines and other putative transmitters between nerve cells, and to the various hormones we have considered in this chapter.

The hormonal systems have been related by physiologists to stress responses and it has been repeatedly suggested that feelings that accompany changes in levels of these hormones are interpreted by the individual according to past learning and present situational cues. Using the conceptual framework of role theory, we may then direct our attention to those points at which stress may be expected and where pattern correlation between physiology and behavior is most likely to be seen within a culture. Conceptualization of these points should be at a level that cuts across specific

roles. In addition to points of stress, there are expected to be physiological relationships to such aspects of roles as dominance or deference relations, and affiliative bonding.

For any individual, the relationship of physiological response to particular role-stress events may be posited as highly predictable. It would seem reasonable, however, when characterizing a population with regard to the relationships of physiology to role stress to think in terms of more general categories. Also, for some individuals, stress repeatedly occurs with interaction in a nurturance system.

The idea of causality in the relations between endocrine and sociological events may be and has been helpful practically as experiments are set up and run. For some purposes the notion of causality is heuristically facilitative. However, if we are indeed dealing with a homeostatic system in terms of the individual's physiology as it adjusts to the psychological and social environment, then overemphasis on causality may, in the broad picture, obscure our vision. But it is not difficult to utilize those conditions on the one hand, while on the other we work toward a more intuitively satisfying understanding of how the individual makes his peace with his world, both physiologically and sociologically.

It is clear that more information, relative to the issue of the manner in

which sociological events interact with physiological variables, will be needed. How do developmental patterns and behavioral states such as status orders or affiliation influence biochemistry? Can the response of endocrine agents or brain neurotransmitter agents to situations later in life be altered by early experience? What types of changes in the various mechanisms involving neuro-regulatory agents can be found in different sociological states? What are the short- and long-term biochemical effects of different types of sociological situations? Does biochemical state influence social behavior? What is the possibility that behavior, such as repeated dominant or submissive behavior, may alter the propensity for long-term or repeated episodes of the behavior? Such studies involve animal investigation, but may lead to human studies. For example, to what extent does a sociological event, e.g., high status or low status, trigger biochemical changes that affect later events, thereby causing a cycle of actions that is neither wholly “sociological” or “psychological,” nor wholly “biochemical?”

From the standpoint of psychiatry, one might well imagine psychiatric illnesses that meet the model of phenylketonuria in which a set of clear-cut, definable biochemical changes leads to severe behavioral changes. Understanding why *those* behavioral changes occur will be a crucial step. On the other hand, there may be other emotional illnesses that are more purely social structural or psychological than in the model just mentioned, and there may be still other illnesses in which a set of social-interaction events or

psychological events in an individual with appropriate biochemical structures lead to an illness.

On the one hand, one encounters those who view mental illness and mind as apart from the brain, and, on the other, those who feel that a twisted thought, a twisted molecule. What is needed is a view that recognizes the subtle interplay of sociological, individual-psychological, biochemical, and physiological processes.

Recent research, including some presented in this paper, suggests the development of a new field of physiological sociology. Such a term is analogous to physiological psychology, and yet recognizes the new approaches and techniques that will be necessary for the field and its concern with behavior related to group structure and processes. The area is exciting in its conception, and yet poses many problems from technical to philosophical. Some of the problems include how to relate biochemical events to sociologically structured events, the effects of drugs on social structures, and developmental processes.

### **Concluding Remarks**

We have chosen to examine some of the endocrine correlates associated with status behavior because status behavior can be viewed as one of the most basic building blocks of structural-social relations. Status structures are



commonly found throughout mammalian species, status behavior has been sufficiently studied, and theories have been constructed, so that it is possible to begin to perform physiological investigations. Studies to date, although few in number with higher primates, suggest the potential of important correlations in a number of endocrine systems, with potential consideration of genetic and developmental processes. Such information can be of importance, theoretically, in terms of understanding aspects of the behavior, and, potentially, in altering some forms of behavior that may be deleterious to the individual. Further, such information may aid in understanding somatic processes in terms of aspects of social structure that profoundly alter somatic function. Such information may prove of value not only in theory, but also, potentially, in the treatment of psychiatric and psychosomatic conditions which would be a particularly important aspect of the new field of physiological sociology.

## Bibliography

- Alexander, B. K. and M. J. Bowers. "The Social Behavior of Japanese Macaques in the Corral," *Primate News*, 6 (1968), 3-10.
- Altmann, S. A. "A Field Study of the Socio biology of Rhesus Monkeys, *Macaca Mulatta*," *Ann. N.Y. Acad. Sci.*, 102 (1962), 338-435.
- Back, K. W. and M. D. Bogdonoff. "Plasma Lipid Responses to Leadership, Conformity, and Deviation," in P. H. Leiderman and D. Shapiro, eds., *Psychobiological Approaches to Social Behavior*, pp. 24-42. 1964. Stanford: Stanford University Press.

- Bales, R. F. "The Equilibrium Problem in Small Groups," in T. Parsons, R. F. Bales, and E. A. Shils, eds., *Working Papers in the Theory of Action*, pp. 111-161. Glencoe, Ill.: Free Press, 1953.
- Bales, R. F. and P. E. Slater. "Role Differentiation," in T. Parsons, R. F. Bales, and J. Olds, eds., *Family Socialization and Interaction Processes*, pp. 259-306. Glencoe, Ill.: Free Press, 1955.
- Bales, R. F., F. L. Strodbeck, T. M. Mills et al. "Channels of Communication in Small Groups," *Am. Soc. Rev.*, 16 (1951) 461-468.
- Barchas, I. Personal communication, 1970.
- Barchas, J. D., R. D. Ciaranello, S. Kessler et al. "Genetic Aspects of the Synthesis of Catecholamines in the Adrenal Medulla," *Psychoneuroendocrinology*. In press.
- . "Genetic Aspects of Catecholamine Synthesis," *Psychoneuroendocrinology*. In press.
- Barchas, J. D., R. D. Ciaranello, J. M. Stolk et al. "Biogenic Amines and Behavior," in S. Levine, ed., *Hormones and Behavior*, pp. 235-329. New York: Academic, 1972.
- Barchas, J. D., J. M. Stolk, R. D. Ciaranello et al. "Neuro-regulatory Agents and Psychological Assessment," in P. McReynolds, ed., *Psychological Assessment*, Vol. 2, pp. 260-292. Palo Alto: Science and Behavior Books, 1971.
- Barchas, J. D. and E. Usdin. *Serotonin and Behavior*. New York: Academic, 1973.
- Barchas, P. R. "Approaches to Aggression as a Social Behavior," in R. Ofshe, ed., *Readings in Social Psychology*, pp. 388-401. Englewood Cliffs, N.J.: Prentice-Hall, 1971.
- . "Differentiation and Stability of Dominance and Deference Orders in Rhesus Monkeys." Ph.D. thesis, Stanford University, 1971. Unpublished.
- Barchas, P. R. and M. H. Fisek. "Status Formation in Ad Hoc Groups of Rhesus and Humans," submitted for publication.
- Barnett, S. A. "Competition among Wild Rats," *Nature*, 175 (1955), 126-127.

- Berger, J. and T. L. Conner. "Performance Expectations and Behavior in Small Groups," *Acta Sociol*, 12 (1969), 186-197.
- Berger, J., B. Cohen, Jr., and M. Zelditch. "Status Characteristics and Expectation States," in J. Berger, M. Zelditch, and B. Anderson, eds., *Sociological Theories in Progress*, Vol. 1, pp. 47-73. Boston: Houghton-Mifflin, 1966.
- Berger, J. and M. H. Fisek. "A Generalization of the Theory of Status Characteristics and Expectation States," in J. Berger, T. L. Connor, and M. H. Fisek, eds., *Expectation States Theory: A Theoretical Research Program*, pp. 163-205. Cambridge, Mass.: Winthrop, 1974.
- Bernstein, I. and W. A. Mason. "Group Formation by Rhesus Monkeys," *Anim. Behav.*, 11 (1963), 28-31.
- Boelkins, R. C. "Effects of Parachlorophenylalanine on the Behavior of Monkeys," in J. D. Barchas and E. Usdin, eds., *Serotonin and Behavior*, pp. 357-364. New York: Academic, 1973.
- Borgatta, E. F. and R. F. Bales. "Interaction of Individuals in Reconstituted Groups," *Sociometry*, 15 (1953), 302-320.
- Brain, P. F. "Endocrine and Behavioral Differences between Dominant and Subordinate Male House Mice Housed in Pairs," *Psychonom. Sci.*, 28 (1972), 260-262.
- Brain, P. F. and N. W. Nowell. "Isolation versus Grouping Effects on Adrenal and Gonadal Function in Albino Mice," *Gen. Comp. Endocrinol.*, 16 (1971), 149-154.
- Carpenter, C. R. *Naturalistic Behavior of Nonhuman Primates*, pp. 342-357; 365-385; 392-397. University Park, Pa.: Pennsylvania State University Press, 1964.
- Champness, B. Quoted in *Time Magazine*, Oct. 17, 1969, p. 74.
- Christian, J. J. "Effect of Population Size on the Weights of the Reproductive Organs of White Mice," *Am. J. Physiol.*, 181 (1955), 477-480.
- Ciaranello, R. D., P. R. Barchas, S. Kessler et al. "Catecholamines: Strain Differences in Biosynthetic

- Enzyme Activity in Mice," *Life Sci.*, 11 (1972), 565-572.
- Ciaranello, R. D., J. N. Dornbusch, and J. D. Barchas. "Regulation of Adrenal Phenylethanolamine N-Methyltransferase Activity in Three Inbred Mouse Strains," *Mol. Pharmacol.*, 8 (1972), 511-520.
- . "Rapid Increase of Phenylethanolamine N-Methyltransferase by Environmental Stress in an Inbred Mouse Strain," *Science*, 175 (1972), 789-790.
- Collias, N. C. "Social Behavior in Animals," *Ecology*, 34 (1953), 810-811.
- Conner, R. L. "Hormones, Biogenic Amines and Aggression," in S. Levine, ed., *Hormones and Behavior*, pp. 209-233. New York: Academic, 1972.
- Darwin, C. *The Expression of the Emotion in Man and Animals*. New York: Appleton, 1896.
- Davis, D. E. and J. J. Christian. "Relation of Adrenal Weight to Social Rank of Mice," *Proc. Soc. Exp. Biol. Med.*, 94 (1957), 728-731.
- Dement, W. C. Personal communication, 1971.
- DeVore, I., ed. *Primate Behavior: Field Studies of Monkeys and Apes*. New York: Holt, Rinehart & Winston, 1965.
- DoAmaral, J. R. Personal communication, 1972.
- Elmadjian, F., J. M. Hope, and E. T. Lamson. "Excretion of Epinephrine in Various Emotional States," *J. Clin. Endocrinol.*, 17 (1957), 608-620.
- Etkin, W. "Theories of Animal Socialization and Communication," in W. Etkin, ed., *Social Behavior and Organization among Vertebrates*, pp. 167-205. Chicago: University of Chicago Press, 1964.
- Fisek, M. H. "The Evolution of Status Structures and Interaction on Task Oriented Discussion Groups." Unpublished Ph.D. dissertation Stanford University, 1968.

- Fisek, M. H. and R. Ofshe. *The Process of Status Evolution*. Technical Report No. 33. Stanford, Calif.: Laboratory for Social Research, Stanford University, 1970.
- Frankenhaeuser, M. "Behavior and Circulating Catecholamines," *Brain Res*, 31 (1971), 241-262.
- Frankenhaeuser, M. and G. Johansson. "Behavior and Catecholamines in Children," in L. Levi, ed., *Society, Stress and Disease, Vol. 1: The Psychosocial Environment and Psychosomatic Diseases*. London: Oxford University Press, forthcoming.
- Goldman, H., L. Lindner, S. Dinitz et al. "The Simple Sociopath: Physiologic and Sociologic Characteristics," *Biol. Psychiatry*, 3 (1971), 77-83.
- Goldstein, M. "Neuroscience Bases of Abnormal Behavior," *Arch. Neurol*, 30 (1974), 1-35.
- Hall, K. R. L. "Aggression in Monkey and Ape Societies," in P. Jay, ed., *Primates: Studies in Adaptation Variability*, pp. 149-161. New York: Holt, Rinehart & Winston, 1968.
- . "Social Training in Monkeys," in P. Jay, ed., *Primates*, pp. 383-397. New York: Holt, Rinehart & Winston, 1968.
- Hamburg, D. A. "Emotions in Perspective of Human Evolution," in P. Knapp, ed., *Expressions of the Emotions in Man*, pp. 300-317. New York: International Universities Press, 1963.
- . "Genetics of Adrenocortical Hormone Metabolism in Relation to Psychological Stress," in J. Hirsch, ed., *Behavior — Genetic Analysis*, pp. 154-175. New York: McGraw-Hill, 1967.
- , ed. *Psychiatry as a Behavioral Science*. Behavioral Social Sciences Survey Monograph Series. Englewood Cliffs, N.J.: Prentice-Hall, 1970.
- Hamburg, D. A., B. Hamburg, and J. Barchas. "Anger and Depression: Current Psychobiological Approaches," in L. Levi, ed., *Parameters of Emotion*. London: Oxford University Press, Forthcoming.
- Hamburg, D. A. and S. Kessler. "A Behavioral-endocrine-genetic Approach to Stress Problems," in S. Pickett, ed., *Memoirs of the Society for Endocrinology No. 15: Endocrine Genetics*, pp. 249-270. London: Cambridge University Press, 1967.

- Hamburg, D. A. and D. Lunde. "Relation of Behavioral, Genetic, and Neuroendocrine Factors to Thyroid Function," in J. Spuhler, ed., *Genetic Diversity and Human Behavior*, pp. 135-170. Chicago: Aldine, 1967.
- Hare, R. D. *Psychopathy: Theory and Research*. New York: Wiley-Interscience, 1970.
- . "Psychopathy and Physiological Responses to Adrenalin," *J. Abnorm. Psychol.*, 79 (1972), 138-147.
- Hare, R. D. and M. J. Quinn. "Psychopathy and Autonomic Conditioning," *J. Abnorm. Psychol.*, 77 (1971), 223-235.
- Hayama, S. "Correlation between Adrenal Gland Weight and Dominance Rank in Caged Crab-eating Monkeys (*Macaca irus*)," *Primates*, 7 (1966), 21-26.
- Hebb, P. O. and W. R. Thompson. "The Social Significance of Animal Studies," in Lindzy, ed., *Handbook of Social Psychology*, Vol. 1, pp. 532-561, New York: Addison-Wesley, 1954.
- Hediger, H. *Wild Animals in Captivity: An Outline of the Biology of Zoological Gardens*. New York: Dover, 1964.
- Heinicke, C. and R. F. Bales. "Developmental Trends on the Structure of Small Groups," *Sociometry*, 16 (1953), 7-38.
- Henry, J. P., D. L. Ely, and P. M. Stephens. "Changes in Catecholamine-controlling Enzymes in Response to Psychosocial Activation of the Defense and Alarm Reactions," in *Physiology, Emotions, and Psychosomatic Illness*, pp. 225-251. London: CIBA Foundation, 1972.
- Henry, J. P., P. M. Stephens, J. Axelrod et al. "Effect of Psychosocial Stimulation on the Enzymes Involved in the Biosynthesis and Metabolism of Noradrenaline and Adrenaline," *Psychosom. Med.*, 33 (1971), 227-237.
- Hinde, R. G. and L. E. Rewell. "Communication by Postures and Facial Expressions in the Rhesus Monkeys (*Macaca mulatta*)," *Proc. Zool. Soc. Inst.*, 138 (1962), 1-21.

- Imanishi, K. "Social Behaviors in Japanese Monkeys, *Macaca fuscata*," *Psychologiozea*, 1 (1957), 47-54.
- . "Social Organization of Subhuman Primates in Their Natural Habitat," *Curr. Anthropol*, 1 (1960), 393-407.
- Jay, P., ed. *Primates: Studies in Adaptation and Variability*. New York: Holt, Rinehart & Winston, 1968.
- Kawai, M. "On the Rank System in a Natural Group of Japanese Monkeys," *Primates*, 1 (1958), 84-98.
- Kessler, S., R. D. Ciaranello, J. G. M. Shire et al. "Genetic Variation in Catecholamine-synthesizing Enzyme Activity," *Proc. Natl. Acad. Sci.*, 69 (1972), 2448-2450.
- Koford, C. B. "Population Dynamics of Rhesus Monkeys on Cayo Santiago," in DeVore, ed., *Primate Behavior*, pp. 160-174. New York: Holt, Rinehart & Winston, 1965.
- Kreuz, L. E., R. M. Rose, and J. R. Jennings. "Suppression of Plasma Testosterone Levels and Psychological Stress," *Arch. Gen. Psychiatry*, 26 (1972), 479-482.
- Leiderman, P. H. and D. Shapiro. *Psychobiological Approaches to Social Behavior*. Stanford, Calif.: Stanford University Press, 1964.
- Leshner, A. I. and D. K. Candland. "Endocrine Effects of Grouping and Dominance Rank in Squirrel Monkeys," *Physiol. Behav.*, 8 (1972), 441-445.
- Levi, I., ed. "Stress and Distress in Response to Psychosocial Stimuli. Laboratory and Real Life Studies on Sympathoadrenomedullary and Related Reactions," *Acta Med. Scand.* (Suppl.), 528 (1972), p. 166.
- Louch, C. D. and M. Higginbotham. "The Relation between Social Rank and Plasma Corticosterone Levels in Mice," *Gen. Comp. Endocrinol.*, 8 (1967), 441-444.
- Maas, J. W., D. E. Redmond, and R. Gauen. "Effects of Serotonin Depletion on Behavior in Monkeys," in J. D. Barchas and E. Usdin, eds., *Serotonin and Behavior*, pp. 351-356.

New York: Academic Press, 1973.

McKinney, W. T., Jr., S. J. Suomi, and F. Harlow. "Depression in Primates," *Am. J. Psychiatry*, 127 (1971), 1313-1320.

Maroney, R. J., J. M. Warren, and M. M. Sinha. "Stability of Social Dominance Hierarchies in Monkeys (*Macaca mulatta*)," *J. Soc. Psychol.*, 50 (1957), 285-293.

Maslow, A. H. "The Role of Dominance in the Social Behavior of Infrahuman Primates : IV. The Determination of Hierarchy in Pairs and in a Group," *J. Genet. Psychol.*, 49 (1936), 161-198.

Mason, J. W. "A Review of Psycho-endocrine Research on the Sympathetic-adrenal Medullary System," *Psychosom. Med.*, 30 (1968), 631-653.

Milkovic, K., C. Winget, T. Deguchi et al. "The Effect of Maternal Manipulation on the Phenylethanolamine N-methyltransferase Activity and Corticosterone Content of the Fetal Adrenal Gland," *Am. J. Physiol.*, 226 (1974), 864-866.

Miller, R. E. and M. V. Murphy. "Social Interactions of Rhesus Monkeys: Food-getting Dominance as a Dependent Variable," *J. Soc. Psychol.*, 44 (1956), 249-255.

Portmann, A. *Animals as Social Beings*. New York: Harper & Row, 1961.

Redmond, D. E., J. W. Maas, A. Kling et al. "Changes in Primate Social Behavior after Treatment with Alpha-Methyl-Paratyrosine," *Psychosom. Med.*, 33 (1971), 97-113.

Redmond, D. E., J. W. Maas, A. Kling et al. "Social Behavior of Monkeys Selectively Depleted of Monoamines," *Science*, 174 (1971), 428-431.

Robins, L. N. *Deviant Children Grown Up: A Sociological and Psychiatric Study of Sociopathic Personality*. Baltimore: Williams & Wilkins, 1966.

Roe, A. and G. G. Simpson, eds. *Behavior and Evolution*. New Haven: Yale University Press, 1958.

Rose, R. M., T. P. Gordon, and I. S. Bernstein. "Plasma Testosterone Levels in the Male Rhesus:



- Influences of Sexual and Social Stimuli," *Science*, 178 (1972), 643-645.
- Rose, R. M., J. W. Holaday, and I. S. Bernstein. "Plasma Testosterone, Dominance Rank and Aggressive Behavior in Male Rhesus Monkeys," *Nature*, 231 (1971), 366-368.
- Rule, C. "A Theory of Human Behavior Based on Studies of Nonhuman Primates," *Perspect. Biol. Med.*, 10 (1967), 153-176.
- Sadleir, R. M. F. S. "The Establishment of a Dominance Rank Order in Male *Peromyscus maniculatus* and Its Stability with Time," *Anim. Behav.*, 18 (1970), 55-59.
- Sassenrath, E. N. "Increased Adrenal Responsiveness Related to Social Stress in Rhesus Monkeys," *Horm. Behav.*, 1 (1969), 283-298.
- Schachter, S. and B. Latané. "Crime, Cognition and the Autonomic Nervous System," in M. R. Jones, ed., *Nebraska Symposium on Motivation, 1964*, pp. 222-272. Lincoln: University of Nebraska Press, 1964.
- Schachter, S. and J. E. Singer. "Cognitive, Social, and Physiological Determinants of Emotional State," *Psychol. Rev.*, 69 (1962), 379-399.
- Schachter, S. and L. Wheeler. "Epinephrine, Chlorpromazine, and Amusement," *J. Abnorm. Soc. Psychol.*, 65 (1962), 121-128.
- Schreier, A. M., H. F. Harlow, and Stollnitz, eds. *Behavior of Non-Human Primates*, Vols. 1 and 2. New York: Academic, 1965.
- Scott, J. P. "Group Formation Determined by Social Behavior: A Comparative Study of Two Mammalian Societies," *Sociometry*, 8 (1945), 42-52.
- . *Animal Behavior*. Chicago: University of Chicago Press, 1958.
- . "The Effects of Early Experience on Social Behavior and Organization," in W. Etkin, ed., *Social Behavior and Organization among Vertebrates*, pp. 231-255. Chicago: University of Chicago Press, 1964.

- Simmel, G. In K. Wolfe, ed., *The Sociology of Georg Simmel*, pp. 181-303. New York: Free Press, 1950.
- Southwick, C. H., ed. *Primate Social Behavior*. Princeton: Van Nostrand, 1963.
- Southwick, C. H., M. A. Beg, and M. R. Siddigi. "A Population Survey of Rhesus Monkeys in Northern India: II. Transportation Routes and First Areas," *Ecology*, 42 (1961), 698-710.
- Stephan, F. and E. G. Mishler. "The Distributions of Participation in Small Groups and Exponential Approximation," *Am. Soc. Rev.*, 17 (1952), 598-608.
- Stolk, J. M., R. Conner, and J. D. Barchas. "Social Environment and Brain Biogenic Amine Metabolism," *J. Comp. Physiol. Psychol.*, in press.
- Terry, R. T. "Primate Grooming as a Tension Reduction Mechanism," *J. Psychol.*, 76 (1970), 129-136.
- Thiessen, D. D. and D. A. Rodgers. "Population Density and Endocrine Function," *Psychol. Bull.*, 58 (1961), 441-451.
- Tinbergen, N. *Social Behavior in Animals, with Special Reference to Vertebrates*. London: Methuen, 1955.
- Tolson, W. W., J. W. Mason, E. J. Sachar et al. "Urinary Catecholamine Responses Associated with Hospital Admission in Normal Human Subjects," *J. Psychosom. Res.*, 8 (1965), 365-372.
- Warren, J. M. and R. J. Maroney. "Competitive Social Interaction between Monkeys," *J. Soc. Psychol.*, 48 (1958), 223-233.
- Welch, B. L. and A. S. Welch. "Greater Lowering of Brain and Adrenal Catecholamines in Group-housed than in Individually-housed Mice Administered DL-a-Methyltyrosine," *J. Pharm. Pharmacol.*, 20 (1968), 244-246.
- Zucherman, S. *The Social Life of Monkeys and Apes*. London: Kegan, Paul, Trench, Trubner, 1932.

## *Notes*

- 1 We should particularly like to thank David Hamburg and Keith Brodie for their encouragement of this chapter. Our work has been graciously supported by the National Institute of Alcohol Abuse and Alcoholism (AA 00498), the National Institute of Mental Health (MH 23861), the Office of Naval Research, and the Grant Foundation. We would like to thank Lynn Hassler, Florence Parma, and Rosemary Schmele for their secretarial assistance.
  
- 2 The reader is referred to Leiderman and Shapiro<sup>71</sup> for a review of experiments that bear on the general subject.